

Running Head: CREATIVITY & FUNCTIONAL CONNECTIVITY

## Creativity at Rest: Exploring Functional Network Connectivity of Creative Experts

William Orwig<sup>a,b</sup>, Roni Setton<sup>a</sup>, Ibai Diez<sup>b</sup>, Elisenda Bueichekú<sup>b</sup>, Megan L. Meyer<sup>c</sup>, Diana I. Tamir<sup>d</sup>, Jorge Sepulcre<sup>b</sup> & Daniel L. Schacter<sup>a</sup>

<sup>a</sup>Harvard University, Department of Psychology, Cambridge, Massachusetts <sup>b</sup>Gordon Center for Medical Imaging, Department of Radiology, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts <sup>c</sup>Dartmouth College, Department of Psychological & Brain Sciences, Hanover, New Hampshire <sup>d</sup>Princeton University, Department of Psychology, Princeton, New Jersey

### Author Note

This research was supported by grants from the National Institutes of Health (R01AG061811 and R01AG061445 to JS). DLS was supported by the National Institute on Aging (AG008441). This research was funded by the Imagination Institute with support from the John Templeton Foundation. The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Correspondence should be addressed to William Orwig, 880 William James Hall, 33 Kirkland Street, Department of Psychology, Harvard University, Cambridge, Massachusetts, 02138; [williamorwig@g.harvard.edu](mailto:williamorwig@g.harvard.edu); ORCID: 0000-0002-9913-6391.

*Keywords:* Creativity; fMRI; Functional Connectivity; Distal Simulation; Vividness

### **Abstract**

The neuroscience of creativity seeks to disentangle the complex brain processes that underpin the generation of novel ideas. Neuroimaging studies of functional connectivity, particularly functional magnetic resonance imaging (fMRI), have revealed individual differences in brain network organization associated with creative ability; however, much of the extant research is limited to laboratory-based divergent thinking measures. To overcome these limitations, we compare functional brain connectivity in a cohort of creative experts (n=29) and controls (n=25) and examine links with creative behavior. First, we replicate prior findings showing reduced connectivity in visual cortex related to higher creative performance. Second, we examine whether this result is driven by integrated or segregated connectivity. Third, we examine associations between functional connectivity and vivid distal simulation separately in creative experts and controls. In accordance with past work, our results show reduced connectivity to the primary visual cortex in creative experts at rest. Additionally, we observe a negative association between distal simulation vividness and connectivity to the lateral visual cortex in creative experts. Taken together, these results highlight connectivity profiles of highly creative people and suggest that creative thinking may be related to, though not fully redundant with, the ability to vividly imagine the future.

## **Creativity at Rest: Exploring Functional Network Connectivity of Creative Experts**

### **1. INTRODUCTION**

The neuroscience of creativity seeks to disentangle the complex brain processes that underpin the generation of novel ideas. Neuroimaging studies of functional connectivity, particularly functional magnetic resonance imaging (fMRI), have revealed individual differences in brain network organization associated with creative ability (Beaty et al., 2014; Takeuchi et al., 2012); however, much of the extant research is limited to laboratory measures of creative performance. Divergent thinking (DT) has been a central concept in creativity research since its first usage by Guilford (1950). The most common assessment of DT is the alternative uses task (AUT), involving the generation of novel uses for common objects. Behavioral studies have shown that DT, as assessed by the AUT, predicts both the quantity of self-reported creative achievements (Jauk, Benedek, & Neubauer, 2014) and the quality of expert-rated creative performance (Beaty et al., 2013), as well as the academic and creative successes of school-aged children (Plucker, 1999, Torrance, 1988). The predictive power of DT has fueled decades of empirical research on the neurocognitive basis of creativity (Cogdell-Brooke et al., 2020; Kim, 2008; Runco & Acar, 2012; Wu et al., 2015). Computerized methods for scoring AUT have led to widely accessible tools for assessing individual differences in creativity (Beaty & Johnson, 2021). While DT certainly captures some aspects of creativity, it is not all-encompassing and its psychometric merit has been the subject of some debate in the field (Forthmann, Jankowska, & Karwowski, 2021; Zeng, Proctor, & Salvendy, 2011). To overcome these limitations, we focus our analyses on a cohort of notable creative experts.

Current theories suggest that creativity is not attributable to a single brain region, rather novel ideas result from interactions between large-scale brain networks (Beaty, Seli, & Schacter,

2019). Among the most well-studied of these functional brain networks, the Default Network (DN) -including midline and posterior inferior parietal regions- shows increased activation in the absence of an external stimulus (Raichle et al., 2001). DN activity has been linked with self-generated thought, such as mind wandering and imagination (Andrews-Hanna et al., 2014; Buckner et al., 2008). Both task-based and resting-state fMRI studies have contributed to current understanding of the neurocognitive basis of creativity. Task-based approaches have helped to uncover the relationship among networks during creative task performance (Beaty et al., 2018; Shi et al., 2018). It has been proposed that DN contributes to the generation of candidate ideas, while executive control networks exert top-down monitoring to meet specific task goals or constraints (Beaty et al., 2014, 2016). Resting-state functional connectivity is thought to reflect experience-dependent patterns of connectivity that relate to individual and group differences in behavior (Stevens & Spreng, 2014). Research on resting-state network organization in relation to creative cognition has investigated inter-individual differences, finding that increased ‘hubness’ (the presence of hubs which form connections between different communities) of DN regions was positively associated with creativity (Kenett et al., 2020). Additionally, a recent study from our group has identified a pattern of reduced resting-state connectivity to the visual cortex associated with increased DT (Orwig et al., 2021). One interpretation of these results is that highly creative people are more prone to engage in internally directed cognition in the absence of an external task. A primary aim of the present study was to replicate this finding of reduced connectivity to visual cortex in a cohort of creative experts.

While most participants in creativity research are drawn from the general population, some recent efforts involve the collection of neuroimaging data from creative professionals. The Big-C Project is a large-scale study, seeking to identify behavioral and functional brain

## CREATIVITY & FUNCTIONAL CONNECTIVITY

characteristics distinguishing exceptionally creative people. For instance, Anderson and colleagues (2022) found that highly creative individuals had higher local clustering coefficients during resting-state and reduced local clustering while performing the AUT compared to less creative individuals. Furthermore, findings from this dataset indicate less activation in bilateral occipital cortex during a DT task in highly creative versus less creative individuals (Japardi et al., 2018). Related work by Chrysikou et al. (2020) described patterns of neural activation between eminent and non-eminent thinkers; despite showing no behavioral differences, eminent thinkers engaged more temporoparietal and less occipital areas than controls during a creative generation task. Additionally, findings from Lotze and colleagues (2014) show reduced interhemispheric connectivity in highly verbally creative individuals compared with less verbally creative individuals.

A second aim is to link creative experts' functional connectivity to a uniquely creative behavior other than DT: vividness of imagination. Imagination refers to the ability to mentally construct and manipulate images that are not directly present to the senses. Imagination is closely associated with creativity, drawing upon semantic (Abraham & Bubic, 2015) and episodic memory processes (Devitt, Addis & Schacter, 2017). Numerous studies conducted in the context of research on episodic memory and future thinking have described the striking cognitive and neural similarities between remembering past experiences and imagining future or other hypothetical experiences (Schacter, Benoit, & Szpunar, 2017; Schacter et al., 2012). Emerging behavioral and neuroimaging research point to a conjunction between episodic retrieval, future imagination, and divergent creative thinking by identifying common engagement of the hippocampus and default network regions (Beaty et al., 2018, Madore et al., 2019; Thakral et al., 2020). Additionally, DT has been linked with the ability to vividly imagine novel and specific

future events (Addis, Pan, Musicaro, & Schacter, 2016; Thakral et al., 2021). Behavioral studies have provided evidence for an association between vividness of mental imagery and divergent thinking (Forisha, 1978; Gonzalez, Campos, & Perez, 1997). Together, these studies highlight the role of constructive episodic processes in creative thinking.

Meyer et al. (2019) examined whether creative experts might be distinguished in their ability to vividly imagine the future. Across two behavioral studies and one neuroimaging study, the authors compared the vividness of *proximal* (i.e., considering what you might do tomorrow) versus *distal* (i.e., considering what life might be like next century) simulation in creative experts and age-matched controls. In both groups, vividness of proximal simulation involved increased medial prefrontal cortex activity. By contrast, creative experts reported having more vivid distal simulations than controls, and showed increased activity of the dorsal medial subnetwork of the DN compared to controls while doing so. The dorsal medial subnetwork includes dorsal medial prefrontal cortex, the temporoparietal junction, along with swaths of inferior frontal and lateral temporal cortex as key structures (Andrews-Hanna et al., 2010) and has been linked to several processes that call on high levels of abstraction, such as semantic processing and mentalizing. Meyer et al. (2019) therefore concluded that an ability to construe abstract thoughts may link creativity and distal simulation. Creative experts in the Meyer et al. (2019) study showed higher connectivity within the dorsal medial DN at rest compared to controls; however, this atlas-based approach is limited, with only the DN tested. This work leaves open the possibility to study whole-brain, voxel-level networks to capture salient features of creative experts' brains.

Building upon our recent work, we revisit the resting-state fMRI data from Meyer et al. (2019), implementing a high resolution, graph theory approach to more precisely characterize global network features of creative thinkers. As with previous analyses (Orwig et al., 2021) a

## CREATIVITY & FUNCTIONAL CONNECTIVITY

voxel-level cortical hubs strategy was used to identify connectivity differences between creative experts and controls without introducing any regional priors. We hypothesize that creative experts will display brain centrality changes in visual areas relative to controls. As creatives showed task-evoked activity differences specific to distal simulation, we also test whether resting-state connectivity may differentially relate to vividness of distal simulation in creatives versus controls. Linking this finding to higher vividness of distal simulations would serve as evidence that highly creative people may more readily silo visual network connectivity and highlight differential patterns of connectivity between creative experts and controls.

## 2. METHODS

### 2.1. Sample

The present research uses behavioral and neuroimaging data previously published in Meyer et al. (2019). We analyze resting-state fMRI data from 53 participants (27 creative experts, 26 controls). Creative experts are defined as any individual who either has been recognized by an award for their creative work, held a position at a prestigious institution known for excellence in a creative domain, or attained commercial success in a creative domain. Specifically, writers, actors, and directors were targeted to capture a range of creative expertise, while still ensuring that creative experts had experience imagining distant times, places, and perspectives. Past research showed that writers and actors/directors demonstrated superior distal simulation skills relative to visual artists; furthermore, creative experts significantly outperformed controls on divergent thinking measures, confirming that ‘real-world’ experts are in fact more creative than controls (Meyer et al., 2019). The sample also includes a control group, targeting professionals working in the legal, medical, and financial industries, based on

## CREATIVITY & FUNCTIONAL CONNECTIVITY

past work that identified individuals in these professions as scoring in the mid-to-low range on standardized creativity assessments (Beketayev & Runco, 2016). Groups did not vary in age (mean creative experts = 36.08 years, SD = 9.85 years; mean controls = 33.73 years, SD = 7.32 years,  $t(50) = 0.98$ ,  $p = .33$ ) or gender ( $\chi^2(1) = 0.03$ ,  $p = 0.88$ ,  $w = 0.001$ ). Participants provided informed consent in accordance with the Princeton University Institutional Review Board (IRB). All materials and data are available on Open Science Framework (<https://osf.io/cy8wt/>).

### 2.2. Distal Simulation Task

Prior to scanning, a subset of participants ( $n = 45$ ; 23 creative experts, 22 controls) completed a distal simulation task, in which they were shown a simulation prompt in the temporal domain (e.g., imagine what the world will be like in 500 years). Participants were shown the prompt for two minutes and instructed to imagine the experience and write a description of their simulation. To derive a subjective measure of vividness, participants then rated the quality of their distal simulations in response to 4 questions: (i) How vividly did you imagine the experience? (ii) To what extent did you see what you imagined in your mind's eye? (iii) To what extent did you feel immersed in what you imagined? and (iv) How difficult was it for you to imagine the experience (reverse scored). Participants responded using a 1 (not at all) to 100 (extremely) sliding scale. A composite score of distal simulation vividness was computed as the average response value across these questions.

### 2.3. MRI acquisition & preprocessing

Scanning was conducted at the Princeton Neuroscience Institute on a 3T Siemens Prisma MRI system with a 64-channel head coil. High resolution T1 scans (MP-RAGE; TR/TE = 2300/2.27, flip angle = 8°, 256 x 256 matrix, 1mm thick, 250mm FoV) were acquired for anatomical normalization. Resting-state functional scans were acquired with a T2\*-weighted



## CREATIVITY & FUNCTIONAL CONNECTIVITY

echo-planar plus sequence with 69 interleaved slices [TR/TE=1500/27ms, flip angle = 75°, 96 x 48 matrix, 2mm thick, 192mm FoV; multiband acceleration factor = 3].

MRI data for both anatomical and functional images were preprocessed using FMRIB Software Library v5.0.7 (FSL) and MATLAB 2017a (Mathworks Inc., Natick, MA). The anatomical and functional preprocessing pipelines were adapted from previous work (Diez et al., 2019). The anatomical T1 preprocessing included: reorientation to right-posterior-inferior (RPI) with `fsloreorient2std`; alignment to anterior and posterior commissures with a custom script; skull stripping using MNI brain mask template projected to individual space; gray matter, white matter and cerebrospinal fluid segmentation using FMRIB's Automated Segmentation Tool (FAST) (Zhang et al., 2001); and computation of non-linear transformation between individual skull-stripped T1 and 2mm resolution MNI152 template images using the FSL `fnirt` tool. The functional MRI preprocessing pipeline included: slice time correction using `Slicetimer`; reorientation to RPI with `fsloreorient2std`; realigning functional volumes within runs with rigid body transformations (6 parameters linear transformation); computation of the transformation between individual skull-stripped T1 and mean functional images using boundary based registration with `flirt`; intensity normalization; removal of confounding factors from the data using linear regression, including 12 motion-related covariates (rigid motion parameters and its derivatives), linear and quadratic terms, and five components each from the lateral ventricles and white matter. Global signal regression was not applied due to the spurious correlations this can introduce (Murphy et al., 2009). Transformation of resting state data to MNI space was performed applying the resulting transformation of concatenating the transformation from functional to structural and from structural to MNI. Spatial smoothing with an isotropic Gaussian kernel of 6-mm FWHM, and band-pass filtering (0.01–0.08 Hz) to reduce low-frequency drift

and high-frequency noise were also applied. Using in-house MATLAB scripts, head motion was quantified using realignment parameters obtained during image preprocessing, including 3 translation and 3 rotation estimates. Scrubbing of time points with excess head motion and interpolation of all time points with Jenkinson framewise displacement  $> 0.2\text{mm}$  was applied. No participants demonstrated excessive head motion; thus, none were removed from the study based on these criteria. The distributions of the correlations across time series were reviewed for possible contamination. No outliers were observed.

### **2.4. Weighted Degree functional connectivity analysis**

Functional connectivity was computed at the individual level using whole-brain voxel-level weighted degree (WD) analysis. WD is a measure of centrality, computed as the sum of the strengths of functional connections between each voxel and the rest of the brain (Bullmore & Sporns, 2009). This centrality measure captures the global features of the networks, which are thought to be relevant in exceptionally creative people (Anderson et al., 2022). WD is one of many possible measures to capture centrality in functional connectivity. Building upon recent findings of WD connectivity associated with individual differences in creativity (Orwig et al., 2021) we applied the same analysis in this context to detect cortical hubs associated with creative expertise. Pearson correlation coefficients were used to calculate the connectivity matrices of each participant using the time series of all cortical gray matter voxels. A r-to-z Fisher transformation was applied to the resulting correlation matrix and negative values were removed due to their controversial interpretation (Qian et al., 2018). To minimize noise, we considered only the most significant links using a false discovery rate (FDR) at  $q\text{-level} < 0.005$  (Benjamini & Hochberg, 1995). After obtaining a high-resolution  $39,080 \times 39,080$  connectivity matrix for each participant, we summed all the weighted connections of each voxel to generate the WD

adjacency matrix. This adjacency matrix was transformed into a brain map and projected on cortical surfaces showing the extent to which each voxel is functionally connected to the rest of the brain.

### **2.5. Integration & Segregation**

Research into the functional composition of network assembly has revealed a modular organization – networks are comprised of densely connected modules, or communities, associated with specific cognitive functions, and cortical hubs which integrate information across communities (Bassett & Bullmore, 2006; Bullmore & Sporns, 2009; Diez et al., 2015). Measures of segregation quantify the presence of communities within the overall brain network, whereas integration captures the brain’s ability to rapidly combine information between communities (Rubinov & Sporns, 2010). To examine group differences in segregation and integration, we organized gray matter voxels with the seventeen-network solution from Yeo and colleagues (2011). If the link’s start and end voxels belonged to the same resting-state network, the link was then classified as a segregated link; otherwise, the link was classified as an integrated link. For each participant, we computed the WD of segregated and integrated links separately obtaining two connectivity maps. Higher values in segregated connectivity maps indicate that a given voxel has a higher number of strong functionally connected links to other voxels within the same functional network, whereas higher values in integrated functional connectivity maps indicate that the voxel is an important hub for integrating information between networks.

### **2.6. Statistical Analysis**

At the behavioral level, we conducted univariate linear regression to test for group differences in distal simulation vividness between creative experts and controls. We report the regression parameters ( $t$  and  $p$  statistics) for this analysis, with a significance threshold of  $\alpha =$

.05. At the neuroimaging level, general linear models were used to compute the group difference between creative experts and controls for whole-brain WD maps, segregation maps, and integration maps. Additionally, general linear models were used to compute the association between WD and distal simulation vividness within creative experts and control groups. Whole-brain correction for multiple comparisons was computed using Monte Carlo simulation with 10,000 iterations to estimate the probability of false positive clusters with a two-tailed  $p$ -value  $< 0.05$  (3dClustSim, afni.nimh.nih.gov). First, we generated the residual of the statistics and used it to estimate the spatial autocorrelation present in our data. Then, we computed null models by generating 10,000 random maps with the estimated spatial autocorrelation (3dClustSim). These null models were used to evaluate the likelihood of getting a brain cluster of a particular size for a  $p$ -value  $< 0.05$  by chance. These data were then used to remove all clusters smaller than the estimated size. Cortical surfaces were visualized using the population-average landmark and surface-based projections of CARET software (Van Essen, 2005). Surface images were displayed using a color scale based on T-scores.

### 3. RESULTS

#### 3.1. Connectivity Patterns Associated with Creative Expertise

We performed whole-brain WD analysis to identify group-level connectivity differences between creative experts and controls. Creative experts showed lower WD connectivity in the medial visual cortex compared to controls (*Figure 1*). Next, we examined whether segregation or integration of networks was driving this effect. Segregation maps did not differ between creative experts and controls; however, integration maps reveal a negative association between connectivity of medial visual cortex and creative expertise. In sum, these results indicate that,

relative to controls, creative experts have reduced connectivity between visual cortex and the rest of the brain during resting-state.

### 3.2. Connectivity Patterns Associated with Distal Simulation Vividness

Behavioral analysis revealed no significant differences between creative experts and controls in relation to vividness of distal simulation ( $t = 1.09$ ,  $p = .28$ ). Given that Meyer et al. (2019) report significant differences in a larger behavioral sample, it is likely that the present sample lacked sufficient statistical power to detect these behavioral differences. To examine the relationship between functional connectivity and distal simulation vividness, we performed two independent linear regression analyses within groups of creative experts and controls. Results indicate a negative association between WD in the lateral visual cortex and distal simulation vividness in creative experts (*Figure 2.a*). Conversely, we find a positive association between WD of voxels in the cingulate cortex and left anterior insula and distal simulation vividness in controls (*Figure 2.b*). These findings reveal differential patterns of connectivity, between creative experts and controls, associated with vividness of distal simulation.

## 4. DISCUSSION

This study aimed to characterize WD connectivity associated with creativity. Our findings indicate that creative experts show reduced WD connectivity of primary visual areas to the rest of the brain at rest compared with controls. This finding extends past work describing individual differences in continuous creative performance within the general population to a group difference present in a highly expert sample of creatives (Orwig et al., 2021). Moreover, we found that reduced WD connectivity to lateral visual cortex was associated with more vivid

## CREATIVITY & FUNCTIONAL CONNECTIVITY

distal simulations in creative experts. Taken together, these findings resemble a similar pattern of connectivity associated with creative expertise and distal imagination.

One interpretation of these findings is that in the absence of an external task –namely, during resting-state– highly creative people are more prone to engage in internally directed cognition, tuning out sensory information from the present environment. Perceptual decoupling, first described by Schooler and colleagues (2011), refers to the capacity to disengage attention from perception during mind wandering. Previous studies of creative cognition have found extended deactivation in occipital cortex associated with internal attention (Benedek et al., 2016). The present findings highlight patterns of reduced WD connectivity to visual cortex in highly creative people. It may be the case that creative experts are engaged in more internally directed cognitive processes during resting-state and this manifests in reduced connectivity to the primary visual areas, compared to controls. An alternative explanation of these results is that involvement in creative pursuits somehow alters the functional organization of resting-state networks, resulting in the observed reduction in connectivity. It is a well-established phenomenon that repeated exposure to the same stimulus results in diminished response in the human visual cortex. Repetition suppression, the relative attenuation of neural signal evoked by repeated presentation of a stimulus, has been investigated extensively using fMRI (Fritsche et al., 2020; Kourtzi & Kanwisher, 2001; Larsson & Smith, 2012). In the context of the present findings, perhaps the observed negative association between creative expertise and visual cortex connectivity is a consequence of training in a creative domain. According to this view, years of experience with writing, acting, or directing may lead to functional changes in the resting-state network organization. These explanations need not be mutually exclusive: it is possible that creative experts engage in more internally-driven thought at rest *and*, in doing so, modify

connectivity of visual cortex. Future work aimed at disentangling the two—either by probing thought content at rest or introducing a novel visual paradigm to examine rates of repetition suppression—will be fruitful in understanding why this functional property is unique to creatives.

Despite the long-standing and intuitive connection between imagination and creative thought, the precise roles of imagery and distal simulation in the creative process remains an open question. While some studies have found a correlation between subjective imagery vividness and creativity (Campos & Pérez, 1989; Shaw & Belmore, 1982), others have failed to observe a reliable association (LeBoutillier & Marks, 2003). Vividness ratings did not differ between the creative experts and controls on which we report here; however, Meyer et al. (2019) did report a difference in their larger behavioral sample. It is likely, then, that the present sample lacked sufficient statistical power. Using this sample, Meyer et al. (2019) reported that creative experts engage regions of the dorsal medial default network during distal simulation more than controls and show higher connectivity within this subnetwork at rest. While Meyer et al. (2019) only tested for group differences within the default network, their results suggested altered brain activity subserving distal simulation across groups that could be detected within functional connectivity patterns at rest. One could argue that failure to observe a behavioral difference in this sub-sample undermines the logic of comparing functional connectivity between creative experts and controls. It is possible, however, that separable neural resources result in comparable behavior. A prime example of this can be seen in the healthy aging literature, where high-performing older adults recruit additional brain resources to reach comparable performance as young adults on working memory tasks (Reuter-Lorenz et al., 2000; Cabeza et al., 2002, 2004; Park & Reuter-Lorenz, 2009). For this reason, it is essential to look at the correlations within the

entire sample as well as separately within each group. Although we acknowledge that the absence of a behavioral difference in our sub-sample limits to some extent the conclusions that we can draw, by describing associations with distal simulation vividness, within groups of creatives and controls, we provide novel insights into the relationship between imagination and creative expertise.

We therefore examined whether vividness would differentially associate with resting-state functional connectivity across groups. We find that vividness for distal future simulation was negatively associated with connectivity to lateral occipital cortex, but only in creative experts. Numerous studies have reported that visual imagery evokes activation in early visual cortex, though others fail to observe this effect (Kosslyn & Thompson, 2003). Here we find that a distinguishing characteristic of the creative brain, namely lower WD connectivity of visual cortex, may support more vivid distal simulation. This is in contrast to healthy controls, wherein higher vividness ratings were associated with greater WD connectivity between regions of the ventral attention network and the rest of the brain. We focused our analysis exclusively on distal simulation, rather than proximal simulation vividness (which was collected with different procedures, while participants were in the scanner) because past work indicates that imagination of distal future events is directly relevant to creativity (Meyer et al., 2019). Although these findings should be replicated in a larger sample, they suggest diverging brain patterns that may be readily recruited for imagination of the far future. If it is the case that creative people more vividly imagine events in the distant future, it could be speculated that, in order to do so, they must attenuate sensory input from their immediate surroundings.

The present research uses resting-state fMRI data to describe WD connectivity of the brain at rest. It should be noted that there is a considerable amount of noise inherent in working



## CREATIVITY & FUNCTIONAL CONNECTIVITY

with resting-state data, given that we are not able to control for participant mood or state of mind during scanning. With these limitations in mind, we utilize graph theory metrics in weighted networks, which have been shown to have robust within-subject reproducibility (Ran et al., 2020). Resting-state functional connectivity is thought to reflect the repeated use of circuits during different tasks and has strong correspondence to task-based connectivity (Stevens & Spreng, 2014). It may be the case that differences in connectivity at rest speak to group differences in how creatives and controls engage in simulation tasks, which may in turn be reflected in resting-state architectures. While future studies may seek to replicate these findings in larger samples of creative experts, this data-driven approach offers initial evidence that connectivity differences at rest may reflect brain-related changes associated with creative expertise. Additionally, assessment of more laboratory measures (e.g., trait-level mind wandering, openness to experience) in future samples of creative experts could advance understanding of the behavioral correlates of creativity. Further analysis of the complex interplay between imagination, vividness and visual cortex promises to offer new insights into the neurocognitive basis of creative thinking.

## REFERENCES

- Abraham, A., & Bubic, A. (2015). Semantic memory as the root of imagination. *Frontiers in psychology, 6*, 325. <https://doi.org/10.3389/fpsyg.2015.00325>
- Addis, D. R., Pan, L., Musicaro, R., & Schacter, D. L. (2016). Divergent thinking and constructing episodic simulations. *Memory (Hove, England), 24*(1), 89–97. <https://doi.org/10.1080/09658211.2014.985591>
- Anderson, A., Japardi, K., Knudsen, K. S., Bookheimer, S. Y., Ghahremani, D. G., & Bilder, R. M. (2022). Big-C creativity in artists and scientists is associated with more random global but less random local fMRI functional connectivity. *Psychology of Aesthetics, Creativity, and the Arts*. Advance online publication. <https://doi.org/10.1037/aca0000463>
- Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-anatomic fractionation of the brain's default network. *Neuron, 65*(4), 550–562. <https://doi.org/10.1016/j.neuron.2010.02.005>
- Bassett, D. S., & Bullmore, E. (2006). Small-world brain networks. *The Neuroscientist: a review journal bringing neurobiology, neurology and psychiatry, 12*(6), 512–523. <https://doi.org/10.1177/1073858406293182>
- Beaty, R. E., Benedek, M., Silvia, P. J., & Schacter, D. L. (2016). Creative cognition and brain network dynamics. *Trends in Cognitive Sciences, 20*(2), 87–95. <https://doi.org/10.1016/j.tics.2015.10.004>
- Beaty, R. E., Benedek, M., Wilkins, R. W., Jauk, E., Fink, A., Silvia, P. J., Hodges, D. A., Koschutnig, K., & Neubauer, A. C. (2014). Creativity and the default network: A

- functional connectivity analysis of the creative brain at rest. *Neuropsychologia*, *64*, 92–98. <https://doi.org/10.1016/j.neuropsychologia.2014.09.019>
- Beaty, R. E., & Johnson, D. R. (2021). Automating creativity assessment with SemDis: An open platform for computing semantic distance. *Behavior research methods*, *53*(2), 757–780. <https://doi.org/10.3758/s13428-020-01453-w>
- Beaty, R. E., Seli, P., & Schacter, D. L. (2019). Network neuroscience of creative cognition: Mapping cognitive mechanisms and individual differences in the creative brain. *Current Opinion in Behavioral Sciences*, *27*, 22–30. <https://doi.org/10.1016/j.cobeha.2018.08.013>
- Beaty, R. E., Smekens, B. A., Silvia, P. J., Hodges, D. A., & Kane, M. J. (2013). A first look at the role of domain-general cognitive and creative abilities in jazz improvisation. *Psychomusicology: Music, Mind, and Brain*, *23*(4), 262–268. <https://doi.org/10.1037/a0034968>
- Beaty, R. E., Thakral, P. P., Madore, K. P., Benedek, M., & Schacter, D. L. (2018). Core network contributions to remembering the past, imagining the future, and thinking creatively. *Journal of Cognitive Neuroscience*, *30*(12), 1939–1951. [https://doi.org/10.1162/jocn\\_a\\_01327](https://doi.org/10.1162/jocn_a_01327)
- Benedek, M., Jauk, E., Beaty, R. E., Fink, A., Koschutnig, K., & Neubauer, A. C. (2016). Brain mechanisms associated with internally directed attention and self-generated thought. *Scientific Reports*, *6*(1), 22959. <https://doi.org/10.1038/srep22959>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>

Beketayev, K., & Runco, M. A. (2016). Scoring Divergent Thinking Tests by Computer With a Semantics-Based Algorithm. *Europe's journal of psychology*, *12*(2), 210–220.

<https://doi.org/10.5964/ejop.v12i2.1127>

Bucci S. M., Maskit B., Bucci, W. (2015). Putting feelings into words: Cross-linguistic markers of the referential process. *Proceedings of the 2nd Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality*, 80-88, Denver, CO.

Bucci W., & Maskit, B. (2006) A weighted referential activity dictionary. *Computing Attitude and Affect in Text: Theory and Applications*, The Information Retrieval Series. (Springer, Dordrecht), pp 49–60.

Bullmore, E., & Sporns, O. (2009). Complex brain networks: Graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience*, *10*(3), 186–198.

<https://doi.org/10.1038/nrn2575>

Cabeza, R., Anderson, N.D., Locantore, J.K., McIntosh, A.R. (2002). Aging gracefully: Compensatory brain activity in high-performing older adults. *Neuroimage*, *17*(3):1394–402.

Cabeza, R., Daselaar, S.M., Dolcos, F., Prince, S.E., Budde, M., Nyberg, L. (2004). Task-independent and task-specific age effects on brain activity during working memory, visual attention and episodic retrieval. *Cerebral Cortex*, *14*(4):364–75.

Campos, A., & González, M. A. (1993). Vividness of imagery and creativity. *Perceptual and Motor Skills*, *77*(3), 923–928. <https://doi.org/10.2466/pms.1993.77.3.923>

Campos, A., & Pérez, M. J. (1989). High and low imagers and their scores on creativity.

*Perceptual and Motor Skills*, 68(2), 403–406. <https://doi.org/10.2466/pms.1989.68.2.403>

Cogdell- Brooke, L. S., Sowden, P. T., Violante, I. R., & Thompson, H. E. (2020). A meta-analysis of functional magnetic resonance imaging studies of divergent thinking using activation likelihood estimation. *Human Brain Mapping*, 41(17), 5057-5077.

<https://doi.org/10.1002/hbm.25170>

Daniels-McGhee, S. & Davis, G. A. (1994). The imagery-creativity connection. *The Journal of Creative Behavior*, 28(3), 151–176. <https://doi.org/10.1002/j.2162-6057.1994.tb01189.x>

Devitt, A. L., Addis, D. R., & Schacter, D. L. (2017). Episodic and semantic content of memory and imagination: A multilevel analysis. *Memory & cognition*, 45(7), 1078–1094.

<https://doi.org/10.3758/s13421-017-0716-1>

Diez, I., Bonifazi, P., Escudero, I., Mateos, B., Muñoz, M. A., Stramaglia, S., & Cortes, J. M. (2015). A novel brain partition highlights the modular skeleton shared by structure and function. *Scientific Reports*, 5, 10532. <https://doi.org/10.1038/srep10532>

Diez, I., Ortiz-Terán, L., Williams, B., Jalilianhasanpour, R., Ospina, J. P., Dickerson, B. C., Keshavan, M. S., Jr, W. C. L., Sepulcre, J., & Perez, D. L. (2019). Corticolimbic fast-tracking: Enhanced multimodal integration in functional neurological disorder. *Journal of Neurology, Neurosurgery & Psychiatry*, 90(8), 929–938. <https://doi.org/10.1136/jnnp-2018-319657>

Feng, Q., He, L., Yang, W., Zhang, Y., Wu, X., & Qiu, J. (2019). Verbal creativity is correlated with the dynamic reconfiguration of brain networks in the resting state. *Frontiers in Psychology*, 10, 894. <https://doi.org/10.3389/fpsyg.2019.00894>

Forisha, B. L. (1978). Mental imagery and creativity: Review and speculations. *Journal of Mental Imagery*, 2(2), 209–238.

Forthmann, B., Jankowska, D. M., & Karwowski, M. (2021). How reliable and valid are frequency-based originality scores? Evidence from a sample of children and adolescents. *Thinking Skills and Creativity*, 41, 100851.

Fritsche, M., Lawrence, S., & de Lange, F. P. (2020). Temporal tuning of repetition suppression across the visual cortex. *Journal of neurophysiology*, 123(1), 224–233.  
<https://doi.org/10.1152/jn.00582.2019>

Gao, Z., Zhang, D., Liang, A., Liang, B., Wang, Z., Cai, Y., Li, J., Gao, M., Liu, X., Chang, S., Jiao, B., Huang, R., & Liu, M. (2017). Exploring the associations between intrinsic brain connectivity and creative ability using functional connectivity strength and connectome analysis. *Brain Connectivity*, 7(9), 590–601. <https://doi.org/10.1089/brain.2017.0510>

Gonzalez, M. A., Campos, A., & Perez, M. J. (1997). Mental imagery and creative thinking. *The Journal of Psychology*, 131(4), 357-364.

Japardi, K., Bookheimer, S., Knudsen, K., Ghahremani, D. G., & Bilder, R. M. (2018). Functional magnetic resonance imaging of divergent and convergent thinking in Big-C creativity. *Neuropsychologia*, 118, 59-67.  
<https://doi.org/10.1016/j.neuropsychologia.2018.02.017>

Jauk, E., Benedek, M., & Neubauer, A. C. (2014). The road to creative achievement: A latent variable model of ability and personality predictors. *European Journal of Personality*, 28(1), 95–105. <https://doi.org/10.1002/per.1941>

- Kim, K. H. (2008). Meta- analyses of the relationship of creative achievement to both IQ and divergent thinking test scores. *The Journal of Creative Behavior*, 42(2), 106-130.
- Kosslyn, S. M., & Thompson, W. L. (2003). When is early visual cortex activated during visual mental imagery? *Psychological bulletin*, 129(5), 723–746. <https://doi.org/10.1037/0033-2909.129.5.723>
- Kourtzi, Z., & Kanwisher, N. (2001). Representation of perceived object shape by the human lateral occipital complex. *Science (New York, N.Y.)*, 293(5534), 1506–1509. <https://doi.org/10.1126/science.1061133>
- Larsson, J., & Smith, A. T. (2012). fMRI repetition suppression: neuronal adaptation or stimulus expectation? *Cerebral cortex (New York, N.Y.: 1991)*, 22(3), 567–576. <https://doi.org/10.1093/cercor/bhr119>
- LeBoutillier, N., & Marks, D. F. (2003). Mental imagery and creativity: a meta-analytic review study. *British journal of psychology (London, England: 1953)*, 94(Pt 1), 29–44. <https://doi.org/10.1348/000712603762842084>
- Li, J., Zhang, D., Liang, A., Liang, B., Wang, Z., Cai, Y., Gao, M., Gao, Z., Chang, S., Jiao, B., Huang, R., & Liu, M. (2017). High transition frequencies of dynamic functional connectivity states in the creative brain. *Scientific Reports*, 7(1), 46072. <https://doi.org/10.1038/srep46072>
- Madore, K. P., Thakral, P. P., Beaty, R. E., Addis, D. R., & Schacter, D. L. (2019). Neural mechanisms of episodic retrieval support divergent creative thinking. *Cerebral Cortex*, 29(1), 150-166. <https://doi.org/10.1093/cercor/bhx312>

## CREATIVITY & FUNCTIONAL CONNECTIVITY

- Meyer, M. L., Hershfield, H. E., Waytz, A. G., Mildner, J. N., & Tamir, D. I. (2019). Creative expertise is associated with transcending the here and now. *Journal of Personality and Social Psychology, 116*(4), 483–494. <https://doi.org/10.1037/pspa0000148>
- Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The impact of global signal regression on resting state correlations: are anti-correlated networks introduced? *NeuroImage, 44*(3), 893–905. <https://doi.org/10.1016/j.neuroimage.2008.09.036>
- Nelson, K. L., Moskowitz, D. J., & Steiner, H. (2008). Narration and vividness as measures of event-specificity in autobiographical memory. *Discourse Processes, 45*(2), 195–209. <https://doi.org/10.1080/01638530701792891>
- Orwig, W., Diez, I., Vannini, P., Beaty, R., & Sepulcre, J. (2021). Creative connections: Computational semantic distance captures individual creativity and resting-state functional connectivity. *Journal of Cognitive Neuroscience, 33*(3), 499–509. [https://doi.org/10.1162/jocn\\_a\\_01658](https://doi.org/10.1162/jocn_a_01658)
- Park, D.C., & Reuter-Lorenz, P. (2009). The adaptive brain: Aging and neurocognitive scaffolding. *Annual Review Psychology, 60*:173-96. <https://doi.org/10.1146/annurev.psych.59.103006.093656>
- Pennebaker, J. (2011). The secret life of pronouns. *New Scientist, 211*(2828), 42-45.
- Plucker, J.A. (1999). Is the proof in the pudding? Reanalyses of Torrance's (1958 to present) longitudinal data. *Creativity Research Journal, 1999; 12*:103–114.



## CREATIVITY & FUNCTIONAL CONNECTIVITY

- Qian, J., Diez, I., Ortiz-Terán, L., Bonadio, C., Liddell, T., Goñi, J., & Sepulcre, J. (2018). Positive connectivity predicts the dynamic intrinsic topology of the human brain network. *Frontiers in Systems Neuroscience, 12*. <https://doi.org/10.3389/fnsys.2018.00038>
- Ran, Q., Jamouille, T., Schaefferbeke, J., Meersmans, K., Vandenberghe, R., & Dupont, P. (2020). Reproducibility of graph measures at the subject level using resting-state fMRI. *Brain and behavior, 10*(8), 2336–2351. <https://doi.org/10.1002/brb3.1705>
- Reuter-Lorenz, P.A., Jonides, J., Smith, E.E., Hartley, A., Miller, A. (2000). Age differences in the frontal lateralization of verbal and spatial working memory revealed by PET. *Journal of Cognitive Neuroscience, 12*(1):174–87.
- Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *NeuroImage, 52*(3), 1059–1069. <https://doi.org/10.1016/j.neuroimage.2009.10.003>
- Runco, M. A., & Acar, S. (2012). Divergent thinking as an indicator of creative potential. *Creativity Research Journal, 24*(1), 66-75.
- Schacter, D. L., Addis, D. R., Hassabis, D., Martin, V. C., Spreng, R. N., & Szpunar, K. K. (2012). The future of memory: remembering, imagining, and the brain. *Neuron, 76*(4), 677-694. <https://doi.org/10.1016/j.neuron.2012.11.001>
- Schacter, D. L., Benoit, R. G., & Szpunar, K. K. (2017). Episodic future thinking: Mechanisms and functions. *Current Opinion in Behavioral Sciences, 17*, 41-50. <https://doi.org/10.1016/j.cobeha.2017.06.002>

- Schooler, J. W., Smallwood, J., Christoff, K., Handy, T. C., Reichle, E. D., & Sayette, M. A. (2011). Meta-awareness, perceptual decoupling and the wandering mind. *Trends in Cognitive Sciences*, S1364661311000878. <https://doi.org/10.1016/j.tics.2011.05.006>
- Shaw, G. A., & Belmore, S. M. (1982). The relationship between imagery and creativity. *Imagination, Cognition and Personality*, 2(2), 115–123. <https://doi.org/10.2190/4RGA-Y1A6-HEK5-LMF8>
- Stevens, W. D., & Spreng, R. N. (2014). Resting-state functional connectivity MRI reveals active processes central to cognition. *Wiley Interdisciplinary Reviews: Cognitive Science*, 5(2), 233–245. <https://doi.org/10.1002/wcs.1275>
- Takeuchi, H., Taki, Y., Hashizume, H., Sassa, Y., Nagase, T., Nouchi, R., & Kawashima, R. (2012). The association between resting functional connectivity and creativity. *Cerebral cortex (New York, N.Y.: 1991)*, 22(12), 2921–2929. <https://doi.org/10.1093/cercor/bhr371>
- Thakral, P.P., Madore, K.P., Kalinowski, S.E., & Schacter, D.L. (2020). Modulation of hippocampal brain networks produces changes in episodic simulation and divergent thinking. *Proceedings of the National Academy of Sciences USA*, 117, 12729-12740. <https://doi.org/10.1073/pnas.2003535117>
- Thakral, P. P., Yang, A. C., Addis, D. R., & Schacter, D. L. (2021). Divergent thinking and constructing future events: dissociating old from new ideas. *Memory (Hove, England)*, 29(6), 729–743. <https://doi.org/10.1080/09658211.2021.1940205>
- Torrance E.P. (1988). In: *The nature of creativity: Contemporary psychological perspectives*. Sternberg R.J., editor. Cambridge University Press; New York: The nature of creativity as manifest in its testing; pp. 43–75.

Van Essen, D. C. (2005). A population-average, landmark- and surface-based (PALS) atlas of human cerebral cortex. *NeuroImage*, *28*(3), 635–662.

<https://doi.org/10.1016/j.neuroimage.2005.06.058>

Wu, X., Yang, W., Tong, D., Sun, J., Chen, Q., Wei, D., Zhang, Q., Zhang, M., & Qiu, J. (2015). A meta-analysis of neuroimaging studies on divergent thinking using activation likelihood estimation. *Human Brain Mapping*, *36*(7), 2703–2718.

<https://doi.org/10.1002/hbm.22801>

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165.

<https://doi.org/10.1152/jn.00338.2011>

Zeng, L., Proctor, R.W., & Salvendy, G. (2011). Can traditional divergent thinking tests be trusted in measuring and predicting real-world creativity? *Creativity Research Journal*, *23*, 24 - 37. <https://doi.org/10.1080/10400419.2011.545713>

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE transactions on medical imaging*, *20*(1), 45–57. <https://doi.org/10.1109/42.906424>


## Figure Legends

### Figure 1. Weighted Degree

WD connectivity across the medial visual cortex was higher in creative expertise compared to controls. Segregation maps did not differ between creative experts and controls; however, integration maps reveal a negative association between connectivity of medial visual cortex and creative expertise.


### Figure 2. Distal Simulation Vividness

(a) Within the group of creative experts, WD of voxels in the lateral visual cortex were negatively associated with distal simulation vividness. (b) Within the control group, WD of voxels in the cingulate cortex and left anterior insula were positively associated with distal simulation vividness.





## Distal Simulation Vividness

### A Creative Experts



### B Controls



### **Author's Summary**

This study sought to uncover the neurocognitive basis of creativity by comparing brain connectivity in a cohort of creative experts and controls. Using graph theory methods, we replicate past work showing reduced connectivity to the primary visual cortex in highly creative people. Additionally, we observed a negative association between vividness of distal future simulation and connectivity to the lateral visual cortex in creative experts. Taken together, these findings highlight connectivity profiles of highly creative people and suggest that creative thinking may be related to the ability to vividly imagine events in the distant future.